arXiv:1210.0652v2 [math.SG] 14 Nov 2012

Dual torus fibrations and homological mirror symmetry for A_n-singularities

Kwokwai Chan and Kazushi Ueda

Abstract

We study homological mirror symmetry for not necessarily compactly supported coherent sheaves on the minimal resolutions of A_n-singularities. An emphasis is put on the relation with the Strominger-Yau-Zaslow conjecture.

1 Introduction

Let Y be the affine hypersurface

$$Y = \left\{ (z, u, v) \in \mathbb{C}^\times \times \mathbb{C}^2 \;\middle|\; uv = z^{-1} f(z) \right\}, \tag{1.1}$$

where $f(z) = (z - a_0)(z - a_1) \cdots (z - a_n)$ is a polynomial of degree $n+1$ with mutually distinct positive real zeros $0 < a_0 < a_1 < \cdots < a_n$. We equip Y with the symplectic structure

$$\omega = -\frac{\sqrt{-1}}{2} \left(\frac{dz \wedge d\bar{z}}{|z|^2} + du \wedge d\overline{u} + dv \wedge d\overline{v} \right)\bigg|_Y .$$

The projection

$$\pi : Y \to \mathbb{C}^\times, \quad (z, u, v) \mapsto z \tag{1.2}$$

is a conic fibration whose discriminant is given by the zeros $\Delta = \{a_0, \dots, a_n\}$ of the polynomial f. Using this, one can show that the map

$$\rho : Y \to B, \quad (z, u, v) \mapsto \left(\log |z|, \frac{1}{2}(|u|^2 - |v|^2) \right) \tag{1.3}$$

is a Lagrangian torus fibration over the base $B = \mathbb{R}^2$, whose discriminant locus is given by

$$\Gamma := \{(s_0, 0), (s_1, 0), \dots, (s_n, 0)\},$$

where $s_i = \log a_i$ for $i = 0, \dots, n$. Each fiber L of this Lagrangian torus fibration is *special* in the sense that one has

$$\mathfrak{Im} \left(e^{\sqrt{-1}\theta} \Omega \right)\Big|_L = 0$$

for some $\theta \in \mathbb{R}$, where

$$\Omega = \operatorname{Res} \frac{dz \wedge du \wedge dv}{zuv - f(z)} = d \log z \wedge d \log u \tag{1.4}$$

is a nowhere-vanishing holomorphic 2-form on Y.

Strominger, Yau, and Zaslow [SYZ96] conjectured that any Calabi-Yau manifold admits a special Lagrangian torus fibration, and its mirror is obtained as the dual torus fibration. In this paper, we apply their ideas on the fibration (1.3), which we will often refer to as the *SYZ fibration*.

Given the SYZ fibration $\rho : Y \to B$, one can equip the complement $B^{\mathrm{sm}} := B \setminus \Gamma$ of the discriminant with two tropical affine structures. One is called the *symplectic affine structure*, and the other is called the *complex affine structure*. Here a manifold with a *tropical affine structure* is a manifold which is obtained by gluing open subsets of $\mathbb{R}^m$ by the action of the affine linear group $\mathbb{R}^m \rtimes GL_m(\mathbb{Z})$ (or $\mathbb{R}^m \rtimes SL_m(\mathbb{Z})$ if the manifold is oriented). An *integral affine manifold* is the special case when the gluing maps belong to $\mathbb{Z}^m \rtimes GL_m(\mathbb{Z})$. The symplectic affine structure is defined by first taking a basis $\{\gamma_1, \gamma_2\}$ of the space of local sections of the relative homology bundle $(R^1\rho_*\mathbb{Z})^\vee$ and integrating the symplectic form ω along these cycles to obtain 1-forms on B; $dx_i = \int_{\gamma_i} \omega$. Local affine coordinate on B^{sm} are primitives of these 1-forms. The complex affine structure is defined similarly by using $\mathfrak{Im}(e^{\sqrt{-1}\theta}\Omega)$ instead of ω, and is *Legendre dual* to the symplectic affine structure [Hit01].

Following earlier works (cf. e.g. [Fuk05, KS06a, GS11, Aur07, Aur09, CLL12, AAK] and references therein), the mirror $\check{Y}$ of Y is identified in [Cha] with the complement of an anti-canonical divisor in the minimal resolution of the A_n-singularity. This mirror $\check{Y}$ admits a special Lagrangian torus fibration, which is an SYZ mirror in the sense that the symplectic and complex affine structures are interchanged between Y and $\check{Y}$.

Let $E_1, \dots, E_n \subset \check{Y}$ be the irreducible components of the exceptional divisor in the minimal resolution. Then there is an isomorphism

$$\begin{array}{cccc} \deg : & \operatorname{Pic} \check{Y} & \to & \mathbb{Z}^n \\ & \Psi & & \Psi \\ & \mathcal{L} & \mapsto & (\deg \mathcal{L}|_{E_i})_{i=1}^n \end{array}$$

of abelian groups. We write the line bundle associated with $\boldsymbol{d} \in \mathbb{Z}^n$ as $\mathcal{L}_{\boldsymbol{d}}$.

Given an SYZ fibration, it is expected that Lagrangian sections of the original manifold and holomorphic line bundles on the mirror manifold are related by a kind of Fourier transform [AP01, LYZ00], which we refer to as the *SYZ transformation*. We introduce the notion of a *strongly admissible path* in $\mathbb{C}^\times \setminus \Delta$, and associate an exact Lagrangian section $L_\gamma \subset Y$ of the SYZ fibration (1.3) to each strongly admissible path. The *winding number* $\boldsymbol{w}(\gamma) = (w_1(\gamma), \dots, w_n(\gamma)) \in \mathbb{Z}^n$ of a strongly admissible path is defined as the intersection numbers with the closed intervals $[a_{i-1}, a_i]$ for $i = 1, \dots, n$.

The first main result in this paper is the following:

Theorem 1.1. *For a strongly admissible path γ, the SYZ transformation of the Lagrangian L_γ is given by the line bundle $\mathcal{L}_{-\boldsymbol{w}(\gamma)}$.*

Next we consider another symplectic manifold defined by

$$Y' = \left\{ (z, u, v) \in \mathbb{C}^\times \times \mathbb{C}^2 \;\middle|\; uv = \frac{1}{z} + z^n \right\}.$$

Note that Y' is a deformation of Y, and hence is symplectomorphic to it. The map (1.3) gives a special Lagrangian torus fibration on Y', whose discriminant consists only of the origin.

The mirror $\check{\mathcal{Y}}$ for Y' is the smooth stack obtained by removing an anti-canonical divisor from the total space $\mathcal{K}$ of the canonical bundle of the weighted projective line $\mathbb{P}(1,n)$. Although the McKay correspondence [KV00] gives a derived equivalence

$$D^b \operatorname{coh} \check{Y} \cong D^b \operatorname{coh} \check{\mathcal{Y}}, \tag{1.5}$$

the Picard groups are *not* isomorphic:

$$\operatorname{Pic} \check{\mathcal{Y}} \cong \operatorname{Pic} \mathbb{P}(n,1) \cong \mathbb{Z} \ncong \mathbb{Z}^n \cong \operatorname{Pic} \check{Y}.$$

Let $\mathcal{O}_{\check{\mathcal{Y}}}(i)$ be the line bundle on $\check{\mathcal{Y}}$ obtained by restricting the pull-back of $\mathcal{O}_{\mathbb{P}(1,n)}(i)$.

Homological mirror symmetry [Kon95, Kon98] for $\mathbb{P}(1,n)$ gives an equivalence

$$D^b \operatorname{coh} \mathbb{P}(1,n) \cong D^b \mathfrak{Fuk}\, W, \tag{1.6}$$

where $\mathfrak{Fuk}\, W$ is a Fukaya category associated with the Lefschetz fibration

$$\begin{array}{cccc} W: & \mathbb{C}^\times & \to & \mathbb{C} \\ & \Uparrow & & \Uparrow \\ & z & \mapsto & 1/z + z^n. \end{array}$$

This is a special case of (a generalization to toric stacks of) the work of Abouzaid [Abo06, Abo09] on homological mirror symmetry for toric varieties. There are also works by Fang [Fan08] and Fang, Liu, Treumann and Zaslow [FLTZ12, FLTZ11] on homological mirror symmetry for toric varieties, which are also motivated by T-duality but different from the work of Abouzaid.

Under the equivalence (1.6), the collection $(\mathcal{O}_{\mathbb{P}(1,n)}(i))_{i=0}^n$ of line bundles is mapped to Lefschetz thimbles $(\Delta_i)_{i=0}^n$. These Lefschetz thimbles can be lifted to Lagrangian sections $(L_i)_{i=0}^n$ of the SYZ fibration for Y'. Let $\mathcal{W}'$ be the full subcategory of the *wrapped Fukaya category* of Y' consisting of $(L_i)_{i=0}^n$. Here the choice for the wrapping Hamiltonian follows that of Pascaleff [Pas].

The second main result in this paper is the following:

Theorem 1.2. *There is an equivalence*

$$D^b \mathcal{W}' \cong D^b \operatorname{coh} \check{\mathcal{Y}} \tag{1.7}$$

of triangulated categories sending L_i *to* $\mathcal{O}_{\check{\mathcal{Y}}}(i)$ *for* $i = 0, \dots, n$.

The proof is based on an analysis of the behavior of the wrapped Fukaya category under suspension, and depends heavily on the work of Pascaleff [Pas]. We expect that $(L_i)_{i=0}^n$ generates the wrapped Fukaya category, so that the left hand side of (1.7) is the whole wrapped Fukaya category. There is also another choice for the wrapping Hamiltonian such that the resulting wrapped Fukaya category is equivalent to $D^b \operatorname{coh} \mathcal{K}$.

Theorems 1.1 and 1.2 are compatible in the following sense: There exists a symplectomorphism $Y \xrightarrow{\sim} Y'$ which induces an equivalence

$$D^b \mathcal{W} \xrightarrow{\sim} D^b \mathcal{W}',$$

so that the Lagrangians $(L_i)_{i=0}^n$ in Y' are images of Lagrangians $(L_{\gamma_i})_{i=0}^n$ in Y associated with certain strongly admissible paths $\gamma_0, \gamma_1, \ldots, \gamma_n$ in $\mathbb{C}^\times \setminus \Delta$. One can then choose a derived equivalence

$$D^b \operatorname{coh} \check{\mathcal{Y}} \xrightarrow{\sim} D^b \operatorname{coh} \check{Y}$$

in such a way that the images of $(L_{\gamma_i})_{i=0}^n$ under the composition

$$D^b \mathcal{W} \xrightarrow{\sim} D^b \mathcal{W}' \xrightarrow{\sim} D^b \operatorname{coh} \check{\mathcal{Y}} \xrightarrow{\sim} D^b \operatorname{coh} \check{Y}$$

of equivalences are precisely given by their SYZ transformations described in Theorem 1.1. This shows that homological mirror symmetry for non-compact branes is realized by SYZ just as for compact branes [Cha].

This paper is organized as follows: In Section 2, we briefly recall the SYZ construction of the mirror manifold $\check{Y}$ from [Cha, Section 3]. In Section 3, we introduce the notion of a strongly admissible path to which we associate a noncompact Lagrangian submanifold in (Y, ω). In Section 4, we describe holomorphic line bundles on $\check{Y}$ obtained as SYZ transformations of noncompact Lagrangian submanifolds associated with strongly admissible paths. In Section 5, we describe a series of conjectures on homological mirror symmetry for a toric Fano stack, which imply homological mirror symmetry for (the complement of an anti-canonical divisor of) the total space of its canonical bundle. In Section 6, we prove slightly weaker versions of these conjectures for the weighted projective line $\mathbb{P}(1, n)$, which are sufficient for the proof of Theorem 1.2.

Acknowledgment: K. C. is supported by Hong Kong RGC Direct Grant for Research 2011/2012 (Project ID: CUHK2060434). K. U. is supported by JSPS Grant-in-Aid for Young Scientists No.24740043. A part of this paper was worked out while K. U. was visiting the Chinese University of Hong Kong, whose hospitality is gratefully acknowledged.

2 SYZ mirror symmetry

We start with the Hamiltonian S^1-action on (Y, ω):

$$e^{2\pi\sqrt{-1}t} \cdot (u, v, z) = (e^{2\pi\sqrt{-1}t} u, e^{-2\pi\sqrt{-1}t} v, z),$$

whose moment map is given by

$$\mu(u, v, z) = \frac{1}{2}(|u|^2 - |v|^2).$$

Let $B := \mathbb{R}^2$. Then the map $\rho : Y \to B$ defined by

$$\rho(u, v, z) = (\log |z|, \mu) = \left(\log |z|, \frac{1}{2}(|u|^2 - |v|^2) \right)$$

is a Lagrangian torus fibration on Y, whose discriminant locus is given by the finite set

$$\Gamma := \{(s_0, 0), (s_1, 0), \ldots, (s_n, 0)\} \subset B,$$

where $s_i := \log a_i$. This is usually called the *SYZ fibration* of (Y, ω).

Let $B^{\mathrm{sm}} := B \backslash \Gamma$ be the smooth locus. Then the fiber $T_{s,\lambda}$ over a point $(s, \lambda) \in B^{\mathrm{sm}}$ is a smooth Lagrangian torus in Y, and each of the fiber $T_{s_i,0}$ over $(s_i, 0) \in \Gamma$ is singular with one nodal singularity. Furthermore, the locus of Lagrangian torus fibers which bound nonconstant holomorphic disks is given by a union of vertical lines:

$$H := \{(s, \lambda) \in B | s = s_i \text{ for some } i = 0, 1, \dots, n\}.$$

Each connected component of H is called a *wall* in B. See [Cha, Proposition 3.1].

The Lagrangian torus fibration $\rho : Y \to B$ induces an integral affine structure on B^{sm}. This is usually called the *symplectic affine structure*. Here, integrality of the affine structure means that the transition maps between charts on B^{sm} are elements of the integral affine linear group

$$\mathrm{Aff}(\mathbb{Z}^2) := \mathrm{GL}_2(\mathbb{Z}) \rtimes \mathbb{Z}^2.$$

This ensures that $\Lambda \subset TB^{\mathrm{sm}}$, the family of lattices locally generated by $\partial/\partial x_1, \partial/\partial x_2$ where x_1, x_2 are local affine coordinates on B^{sm}, is well-defined.

In 1996, Strominger, Yau and Zaslow [SYZ96] proposed that the mirror of (Y, ω) should be given by fiberwise-dualizing the SYZ fibration on (Y, ω). More precisely, one defines the *semi-flat mirror* of (Y, ω) as the moduli space $\check{Y}_0$ of pairs $(T_{s,\lambda}, \nabla)$ where ∇ is a flat $U(1)$-connection (up to gauge equivalence) on the trivial line bundle $\underline{\mathbb{C}} := \mathbb{C} \times T_{s,\lambda}$ over a smooth Lagrangian torus fiber $T_{s,\lambda}$. Topologically, $\check{Y}_0$ is the quotient TB^{sm}/Λ of the tangent bundle TB^{sm} by Λ, and hence is naturally a complex manifold where the local complex coordinates are given by exponentiations of complexifications of local affine coordinates on B^{sm} (while the quotient $T^*B^{\mathrm{sm}}/\Lambda^\vee$ of the cotangent bundle T^*B^{sm} by the dual lattice $\Lambda^\vee$, which is canonically a symplectic manifold, is contained in Y as an open dense subset such that the restriction of ω to it gives the canonical symplectic structure). However, this is not quite the correct mirror manifold because the natural complex structure on $\check{Y}_0$ is not globally defined due to nontrivial monodromy of the affine structure around each singular point $(s_i, 0) \in \Gamma$.

A more concrete description of this phenomenon is as follows. For $i = 0, 1, \dots, n$, consider the strip $B_i := (s_{i-1}, s_{i+1}) \times \mathbb{R}$ (where we set $s_{-1} := -\infty$ and $s_{n+1} := +\infty$). A covering of B^{sm} is given by the open sets

$$U_i := B_i \setminus [s_i, s_{i+1}) \times \{0\},\ V_i := B_i \setminus (s_{i-1}, s_i] \times \{0\}.$$

$i = 0, 1, \dots, n$. The intersection $U_i \cap V_i$ consists of two connected components:

$$U_i \cap V_i = B_i^+ \cup B_i^-,$$

where B_i^+ (resp. B_i^-) corresponds to the component in which $\lambda > 0$ (resp. $\lambda < 0$).

On U_i (resp. V_i), denote by u_i (resp. v_{i+1}) the exponentiation of the complexification of the left-pointing (resp. right-pointing) affine coordinate. (Our convention is that for a real number $x \in \mathbb{R}$, its complexification is given by $-x + \sqrt{-1}y$.) Also, denote by w the exponentiation of the complexification of the upward-pointing affine coordinate. See Figure 2.1.

Now w is a globally defined coordinate. Geometrically, it can be written as

$$w(T_{s,\lambda}, \nabla) = \begin{cases} \exp(-\int_\alpha \omega)\mathrm{hol}_\nabla(\partial\alpha) & \text{for } \lambda \geq 0, \\ \exp(\int_\alpha \omega)\mathrm{hol}_\nabla(-\partial\alpha) & \text{for } \lambda \leq 0, \end{cases}$$



Figure 2.1: The base affine manifold B.

where $\alpha \in \pi_2(Y, T_{s_i,\lambda})$ is the class of the holomorphic disk in Y bounded by $T_{s_i,\lambda}$ for some $i = 0, 1, \ldots, n$, and $\mathrm{hol}_\nabla(\partial\alpha)$ is the holonomy of the flat $U(1)$-connection ∇ around the boundary $\partial\alpha \in \pi_1(T_{s_i,\lambda})$. The class α changes to $-\alpha$ when one moves from $\lambda > 0$ to $\lambda < 0$.

But the other coordinates u_i and v_{i+1} are not globally defined: Since the monodromy of the affine structure going counter-clockwise around $(s_i, 0) \in \Gamma$ is given by the matrix

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix},$$

the coordinates (u_i, w) and (v_{i+1}, w) glue on $U_i \cap V_i$ according to

$$\begin{aligned} u_i &= v_{i+1}^{-1} \quad \text{on } B_i^+, \\ u_i &= v_{i+1}^{-1} w \quad \text{on } B_i^-. \end{aligned}$$

Hence, the monodromy of the complex coordinates going counter-clockwise around the point $(s_i, 0)$ is nontrivial and given by

$$u_i \mapsto u_i w, \; v_{i+1}^{-1} \mapsto v_{i+1}^{-1} w.$$

In particular, these complex coordinates on TB_i/Λ do not form a globally defined complex structure on $\check{Y}_0$.

The examples in [Aur07, Aur09, Cha10, CLL12] suggest the following construction (we should mention that these are all special cases of the constructions in Kontsevich-Soibelman [KS06b] and Gross-Siebert [GS11], but those general constructions largely ignore the symplectic aspects and hence Lagrangian torus fibration structures - they build the mirror family directly from tropical data in the base B): in order to get the genuine mirror manifold, we need to modify the gluing of complex charts on $\check{Y}_0$ by quantum corrections from disk instantons

$$u_i = v_{i+1}^{-1} + v_{i+1}^{-1} w = \quad v_{i+1}^{-1}(1 + w) \quad \text{on } B_i^+, \tag{2.1}$$

$$u_i = v_{i+1}^{-1} w + v_{i+1}^{-1} = \quad v_{i+1}^{-1} w(1 + w^{-1}) \quad \text{on } B_i^-. \tag{2.2}$$

Geometrically, the term $v_{i+1}^{-1}w$ in the first formula (2.1) should be viewed as multiplying v_{i+1}^{-1} by w where w corresponds to the unique nonconstant holomorphic disk bounded by the Lagrangian torus $T_{0,\lambda}$ whose area is given by

$$\lambda = -\log|w| > 0.$$

This means that we are correcting the term v_{i+1}^{-1} by adding the contribution from the holomorphic disk bounded by $T_{0,\lambda}$ when we cross the upper half of the wall $\{s_i\} \times \mathbb{R} \subset H$. In the same way, the term v_{i+1}^{-1} in the second formula (2.2) is given by multiplying $v_{i+1}^{-1}w$ by w^{-1} where w^{-1} now corresponds to the unique nonconstant holomorphic disk bounded by the Lagrangian torus $T_{0,\lambda}$ whose area is equal to

$$-\lambda = \log|w| = -\log|w^{-1}| > 0.$$

So we are correcting $v_{i+1}^{-1}w$ by the disk bounded by $T_{0,\lambda}$ when we cross the lower half of the wall $\{s_i\} \times \mathbb{R} \subset H$.

The formulas (2.1), (2.2) can actually be interpreted as wall-crossing formulas for the counting of Maslov index two holomorphic disks in Y bounded by Lagrangian torus fibers. To see this, we need to partially compactify Y by allowing z to take values in $\mathbb{C}$ and replacing ρ by the map

$$(u, v, z) \mapsto (|z|, \mu) = (|z|, (|u|^2 - |v|^2)/2).$$

Then the base becomes an affine manifold with both singularities and boundary: it is a right half-space $\bar{B}$ in $\mathbb{R}^2$, and the boundary $\partial\bar{B}$ corresponds to the divisor in Y defined by $z = 0$. In this situation, each of the local coordinates u_i, v_{i+1}^{-1} and $v_{i+1}^{-1}w$ can be expressed as in the form

$$\exp\left(-\int_\beta \omega\right)\mathrm{hol}_\nabla(\partial\beta),$$

for a suitable relative homotopy class $\beta \in \pi_2(Y, T_{s,\lambda})$ of Maslov index two (cf. Auroux [Aur07, Aur09]).

In any case, the modification of gluing cancels the monodromy and defines global complex coordinates u_i, v_i, w on TB_i/Λ (topologically, $TB_i/\Lambda \cong (\mathbb{C}^\times)^2$) related by

$$u_i = v_i^{-1},\ u_{i+1} = v_{i+1}^{-1} \text{ and } u_i v_{i+1} = 1 + w. \tag{2.3}$$

The instanton-corrected mirror $\check{Y}$ is then obtained by gluing together the pieces TB_i/Λ ($i = 0, 1, \ldots, n$) according to (2.3). On the intersection $(TB_{i-1}/\Lambda) \cap (TB_i/\Lambda)$, we have

$$u_i = v_i^{-1},\ u_i v_{i+1} = 1 + w = u_{i-1}v_i.$$

We remark that, by our definition, the mirror manifold $\check{Y}$ will have "gaps" because for instance u, v, w are $\mathbb{C}^\times$-valued. A natural way to "complete" the mirror manifold and fill out its gaps is by rescaling the symplectic structure of Y and performing analytic continuation; see [Aur07, Section 4.2] for a discussion of this "renormalization" process.

Finally, one observes that this is precisely the gluing of complex charts in the toric resolution $X \to \mathbb{C}^2/\mathbb{Z}_{n+1}$ of the A_n-singularity. We thus conclude that the mirror is precisely given by the complex manifold

$$\check{Y} := X \setminus D,$$

where $X = X_\Sigma$ is the toric surface defined by the 2-dimensional fan $\Sigma \in \mathbb{R}^2$ generated by

$$\{v_i := (i, 1) \in N | i = -1, 0, 1, \ldots, n\},$$

and D is the hypersurface in X_Σ defined by $h := \chi^{(0,1)} = 1$ (here $h : X \to \mathbb{C}$ is the holomorphic function whose zero locus is the union of all the toric divisors in X). For $i = 1, \ldots, n$, if we denote by S_i the interval $(s_{i-1}, s_i) \times \{0\} \subset B$, then (the closure of) the quotient $TS_i/TS_i \cap \Lambda$ in $\check{Y}$ is one of the n exceptional divisors (each is a (-2)-curve) in X_Σ. We denote this exceptional divisor by $E_i \subset \check{Y}$.

3 Lagrangian submanifolds fibred over paths

Consider the projection map

$$\pi : Y \to \mathbb{C}^\times, \ (u, v, z) \mapsto z.$$

Each fiber is a conic $\{(u, v) \in \mathbb{C}^2 | uv = z^{-1} f(z)\}$ which degenerates to the cone $uv = 0$ over a zero of the polynomial f. Let us denote by

$$\Delta := \{a_0, a_1, \ldots, a_n\}$$

the set of zeros of the polynomial $f(z)$. Recall that these all real and positive such that

$$0 < a_0 < a_1 < \ldots < a_n.$$

The symplectic fibration $\pi : Y \to \mathbb{C}^\times$ induces a connection on the tangent bundle of Y where the horizontal subspace at $y \in Y_0$ is given by the symplectic orthogonal complement to the vertical subspace:

$$H_y := \mathrm{Ker}(d\pi_y)^{\perp_\omega}.$$

Alternatively, since (Y, ω) is Kähler with respect to the natural complex structure on Y, the horizontal subspace can be regarded as the orthogonal complement to $\mathrm{Ker}(d\pi_y)$ with respect to the Kähler metric.

Given a smooth embedded path $\gamma : I \to \mathbb{C}^\times \setminus \Delta$ where $I \subset \mathbb{R}$ is an interval, parallel transport with respect to the above connection along γ yields symplectomorphisms between smooth fibers

$$\tau_{t_1}^{t_2} : \pi^{-1}(\gamma(t_1)) \to \pi^{-1}(\gamma(t_2)),$$

for $t_1 < t_2 \in I$. These symplectomorphisms are S^1-equivariant since the S^1-action is Hamiltonian. The vanishing cycle V_t in a smooth fiber $\pi^{-1}(\gamma(t))$ is given by its equator $\mu = 0$ (i.e. $|u| = |v|$). Vanishing cycles are invariant under parallel transport, so that the *Lefschetz thimble* L_γ along γ, given by the union of all vanishing cycles $V_t \subset \pi^{-1}(\gamma(t))$ for $t \in I$, is a Lagrangian submanifold in (Y, ω). If γ is a matching path, i.e. a path connecting two critical values of π in $\mathbb{C}^\times$, then L_γ is a Lagrangian sphere in (Y, ω).

Other Lagrangian submanifolds can be constructed in a similar way, as has been done in the recent work of Pascaleff [Pas].

Definition 3.1. Let $\gamma : \mathbb{R} \to \mathbb{C}^\times \setminus \Delta$ be a smooth embedded path. We call the path γ *admissible* if it satisfies the following conditions:

(1) $\gamma(t)$ lies in the negative real axis whenever $|t|$ is sufficiently large, i.e. there exists $T > 0$ such that $\gamma(t) \in \mathbb{R}_{<0}$ if $t > T$ or $t < -T$.

(2) $\lim_{t\to-\infty} \gamma(t) = 0$ and $\lim_{t\to\infty} \gamma(t) = -\infty$.

Given an admissible path $\gamma : \mathbb{R} \to \mathbb{C}^\times$, we consider the Lagrangian (real locus)

$$C_t := \{(u, v, z) \in \pi^{-1}(\gamma(t)) | u, v \in \mathbb{R}\}.$$

in the conic fiber $\pi^{-1}(\gamma(t))$ for each $t < -T$ (see Figure 5.5), which is 'dual' to the vanishing cycle V_t. Notice that C_t is invariant under symplectic parallel transport for all $t < -T$; more precisely, this means that

$$\tau_{t_1}^{t_2}(C_{t_1}) = C_{t_2}$$

for all $t_1 < t_2 < -T$. Let L_γ be the submanifold in Y swept out by parallel transport of C_t $(t < -T)$:

$$L_\gamma := \bigcup_{t\in(-\infty,-T)} C_t \cup \bigcup_{t\in[-T,\infty)} \tau_{-T}^{t}(C_{-T}).$$

This defines a Lagrangian submanifold in (Y, ω) which is homeomorphic to $\mathbb{R}^2$. Note that the curve $C_{-T} := \{(u, v, z) \in \pi^{-1}(\gamma(-T)) | u, v \in \mathbb{R}\}$ in the conic fiber $\pi^{-1}(\gamma(-T))$ is being twisted by Dehn twists in vanishing cycles (i.e. the equator $\lambda = 0$ in a fiber of $\pi : Y \to \mathbb{C}^\times$) as one goes along γ.

Definition 3.2. Let $\gamma : \mathbb{R} \to \mathbb{C}^\times \setminus \Delta$ be an admissible path. For $i = 1, \dots, n$, we define the *i-th winding number* $w_i(\gamma)$ of γ to be the topological intersection number between $\gamma(\mathbb{R})$ and the line segment $[s_{i-1}, s_i] \subset \mathbb{C}^\times$.

Notice that the Hamiltonian isotopy class of the Lagrangian submanifold L_γ and the winding numbers of γ are invariant when we deform γ in a fixed homotopy class (relative to the ends $\gamma|_{(-\infty,-T]}$ and $\gamma|_{[T,\infty]}$). In particular, we can deform γ so that $\gamma(s_i) = -a_i$ for $0 \le i \le n$ (up to a re-parametrization if necessary). In this case, the winding numbers can be computed as follows: There exists a unique lift $\tilde{\gamma} : [s_0, \infty) \to \mathbb{R}$ of $\gamma/|\gamma| : [s_0, \infty) \to S^1 = \mathbb{R}/2\pi\mathbb{Z}$ such that $\tilde{\gamma}(s_0) = 0$. Then for $i = 1, \dots, n$, the i-th winding number of γ is given by the integer

$$w_i(\gamma) = \frac{1}{2\pi}(\tilde{\gamma}(s_i) - \tilde{\gamma}(s_{i-1})).$$

Furthermore, given an admissible path γ, we can in fact deform it (again in a fixed homotopy class fixing the ends $\gamma|_{(-\infty,-T]}$ and $\gamma|_{[T,\infty]}$) so that the following condition is satisfied:

Definition 3.3. Let $\gamma : \mathbb{R} \to \mathbb{C}^\times \setminus \Delta$ be an admissible path. We call γ *strongly admissible* if γ intersects each circle in $\mathbb{C}^\times$ centered at the origin once, i.e. $|\gamma(\mathbb{R}) \cap \{z \in \mathbb{C}^\times | |z| = e^s\}| = 1$ for all $s \in \mathbb{R}$, or equivalently, if the modulus $|\gamma(t)|$ is strictly increasing.

This condition ensures that the noncompact Lagrangian submanifold L_γ is a section of the SYZ fibration $\rho : Y \to B$ (see Section 2):

Proposition 3.4. *Let* $\gamma : \mathbb{R} \to \mathbb{C}^\times$ *be a strongly admissible path. Then the Lagrangian submanifold* L_γ *is a section of the SYZ fibration* $\rho : Y \to B$.

Proof. By definition, L_γ is invariant under parallel transport, so it is a Lagrangian submanifold in (Y, ω). Moreover, it is clear that the moment map μ when restricted to C_t for $t < -T$ is a one-to-one map. Since the symplectomorphisms τ^t_{-T} are all S^1-equivariant, the value of μ is preserved under parallel transport. Thus the restriction of μ to $\tau^t_{-T}(C_{-T})$ remains a one-to-one map. Together with the condition that $|\gamma(t)|$ is strictly increasing, this implies that L_γ intersects each fiber $T_{s,\lambda}$ of the SYZ fibration $\rho : Y \to B$ at one point. Hence L_γ is a section of $\rho : Y \to B$. $\square$

As an example, consider the path

$$\gamma_0 : \mathbb{R} \to \mathbb{C}^\times,\ t \mapsto -e^t,$$

which runs through the whole negative real axis. This is obviously strongly admissible. The corresponding Lagrangian submanifold $L_0 := L_{\gamma_0}$ is simply the Cartesian product of the curve C_{s_0} inside the conic fiber $C_{s_0} = \pi^{-1}(-a_0)$ with the negative real axis (which is the image $\gamma_0(\mathbb{R})$ of the path). As a section of the SYZ fibration $\rho : Y \to B$, L_0 can be explicitly written as

$$\sigma : B \to Y,\ (s, \lambda) \mapsto (u(s,\lambda), v(s,\lambda), z(s,\lambda)),$$

where

$$\begin{aligned} u(s,\lambda) &= \sqrt{\sqrt{\lambda^2 + |f(-e^s)|^2} + \lambda}, \\ v(s,\lambda) &= (-1)^{n-1}\sqrt{\sqrt{\lambda^2 + |f(-e^s)|^2} - \lambda}, \\ z(s,\lambda) &= -e^s. \end{aligned}$$

4 SYZ transformations

Consider $\Lambda^\vee \subset T^*B^{\text{sm}}$, the family of lattices locally generated by dx_1, dx_2. As before, x_1, x_2 here denote local affine coordinates on B^{sm}. As in [Cha], we will assume that $x_2 = -\lambda$ is the globally defined coordinate. Let

$$\omega_0 := dx_1 \wedge d\xi_1 + dx_2 \wedge d\xi_2$$

be the canonical symplectic structure on the quotient $T^*B^{\text{sm}}/\Lambda^\vee$ of the cotangent bundle T^*B^{sm} by $\Lambda^\vee$, where (ξ_1, ξ_2) are fiber coordinates on T^*B^{sm} so that $(x_1, x_2, \xi_1, \xi_2) \in T^*B^{\text{sm}}$ denotes the cotangent vector $\xi_1 dx_1 + \xi_2 dx_2$ at the point $(x_1, x_2) \in B^{\text{sm}}$. In the previous section, we have constructed a global Lagrangian section L_0 of the SYZ fibration $\rho : Y \to B$. Then a theorem of Duistermaat [Dui80] says that there exists a fiber-preserving symplectomorphism

$$\Theta : (T^*B^{\text{sm}}/\Lambda^\vee, \omega_0) \xrightarrow{\cong} (\rho^{-1}(B^{\text{sm}}), \omega)$$

so that L_0 corresponds to the zero section of $T^*B^{\text{sm}}/\Lambda^\vee$.

This symplectomorphism can be constructed as follows. Let $b \in B^{\mathrm{sm}}$. Then for every $\alpha \in T_b^* B^{\mathrm{sm}}$, we can associate a vector field v_α on the fiber $\rho^{-1}(b)$ by

$$\iota_{v_\alpha}\omega = \rho^*\alpha.$$

Let ϕ_α^τ be the flow of v_α at time $\tau \in \mathbb{R}$. Then we define an action θ_α of α on $\rho^{-1}(b)$ by the time-1 flow

$$\theta_\alpha(y) = \phi_\alpha^1(y),\ y \in \rho^{-1}(b).$$

Now the covering map

$$T^*B^{\mathrm{sm}} \to \rho^{-1}(B^{\mathrm{sm}}),\ \alpha \mapsto \theta_\alpha(\sigma(\mathrm{pr}(\alpha))),$$

where $\mathrm{pr} : T^*B^{\mathrm{sm}} \to B^{\mathrm{sm}}$ denotes the projection map, induces a symplectomorphism $\Theta : T^*B^{\mathrm{sm}}/\Lambda^\vee \to \rho^{-1}(B^{\mathrm{sm}})$.

Now let y_1, y_2 be the fiber coordinates on TB^{sm} dual to ξ_1, ξ_2, i.e. $(x_1, x_2, y_1, y_2) \in TB^{\mathrm{sm}}$ denotes the tangent vector $y_1\partial/\partial x_1 + y_2\partial/\partial x_2$ at the point $(x_1, x_2) \in B^{\mathrm{sm}}$. Given a strongly admissible path $\gamma : \mathbb{R} \to \mathbb{C}^\times$, the noncompact Lagrangian submanifold L_γ is a section of the SYZ fibration $\rho : Y \to B$ by Proposition 3.4 (and as we mentioned before, every admissible path can be deformed to a strongly admissible one). Via the symplectomorphism Θ, we can write L_γ in the form

$$L_\gamma = \{(x_1, x_2, \xi_1, \xi_2) \in T^*B^{\mathrm{sm}}/\Lambda^\vee | (x_1, x_2) \in B^{\mathrm{sm}},\ \xi_j = \xi_j(x_1, x_2) \text{ for } j = 1, 2\},$$

where $\xi_j = \xi_j(x_1, x_2)$ $(j = 1, 2)$ are smooth functions on B^{sm}. Since L_γ is Lagrangian, the functions ξ_1, ξ_2 satisfy

$$\frac{\partial \xi_j}{\partial x_l} = \frac{\partial \xi_l}{\partial x_j}$$

for $j, l = 1, 2$ (see [LYZ00, AP01, Cha]).

Lying at the basis of the SYZ proposal [SYZ96] is the fact that the dual T^* of a torus T can be viewed as the moduli space of flat $U(1)$-connections on the trivial line bundle $\underline{\mathbb{C}} := \mathbb{C} \times T$ over T. So a Lagrangian section $L = \{(x, \xi(x)) \in T^*U/T^*U \cap \Lambda^\vee | x \in U\}$ over an open set $U \subset B^{\mathrm{sm}}$ corresponds to a family of connections $\{\nabla_{\xi(x)} | x \in U\}$ patching together to give a $U(1)$-connection which can locally be written as

$$\check{\nabla}_U = d + 2\pi\sqrt{-1}(\xi_1 dy_1 + \xi_2 dy_2)$$

over $TU/TU \cap \Lambda \subset \check{Y}$. As shown in [LYZ00, AP01] (see also [Cha, Section 2]), the $(0, 2)$-part $F^{(0,2)}$ (and also $(2, 0)$-part $F^{(2,0)}$) of the curvature two form of $\check{\nabla}_U$ is trivial.

Recall that a covering of B^{sm} is given by the open sets

$$U_i := B_i \setminus [s_i, s_{i+1}) \times \{0\},\ V_i := B_i \setminus (s_{i-1}, s_i] \times \{0\}.$$

for $i = 0, 1, \ldots, n$. Now, the restriction of the Lagrangian section L_γ to each U_i (resp. V_i) is transformed to a $U(1)$-connection $\check{\nabla}_{U_i}$ over $TU_i/TU_i \cap \Lambda$ (resp. $\check{\nabla}_{V_i}$ over $TV_i/TV_i \cap \Lambda$). These connections can be glued together according to the gluing formulas (2.1), (2.2). Since the $(0, 2)$-part $F^{(0,2)}$ of the curvature two form vanishes, this defines a holomorphic line bundle $\mathcal{L}_\gamma$ over $\check{Y}$.

Definition 4.1. Let $\gamma : \mathbb{R} \to \mathbb{C}^\times$ be a strongly admissible path and L_γ be the noncompact Lagrangian submanifold in (Y, ω) associated to γ. We define the SYZ transformation of L_γ to be the holomorphic line bundle $\mathcal{L}_\gamma$ over $\check{Y}$, i.e.

$$\mathcal{F}(L_\gamma) := \mathcal{L}_\gamma.$$

Notice that the isomorphism class of $\mathcal{L}_\gamma$ is unchanged when we deform L_γ in a fixed Hamiltonian isotopy class (or deforming γ is a fixed homotopy class relative to the ends of γ). Henceforth, we will regard this as defining the SYZ transformation of the Hamiltonian isotopy class of the Lagrangian submanifold $L_\gamma \subset Y$ as an isomorphism class of holomorphic line bundle over $\check{Y}$.

As an immediate example, the SYZ transformation of the zero section $L_0 = L_{\gamma_0}$ gives the structure sheaf $\mathcal{O}_{\check{Y}}$ over $\check{Y}$.

The main goal of this section is to compute (the isomorphism class of) the line bundle $\mathcal{L}_\gamma$ in terms of the winding numbers of the path γ. To begin with, note that the isomorphism class of a line bundle over $\check{Y}$ is determined by the degrees of its restrictions to the exceptional divisors E_i for $i = 1, \ldots, n$. Given integers $d_1, \ldots, d_n \in \mathbb{Z}$, let us denote by $\mathcal{L}_{d_1,\ldots,d_n}$ the line bundle on $\check{Y}$ such that $\deg \mathcal{L}_{d_1,\ldots,d_n}|_{E_i} = d_i$.

Now, given a Lagrangian section

$$L_\gamma = \{(x_1, x_2, \xi_1, \xi_2) \in T^*B^{\mathrm{sm}}/\Lambda^\vee | (x_1, x_2) \in B^{\mathrm{sm}},\ \xi_j = \xi_j(x_1, x_2) \text{ for } j = 1, 2\},$$

of the SYZ fibration $\rho : Y \to B$, its SYZ transformation is the connection $\check{\nabla}$ which can locally be expressed as

$$\check{\nabla}_U = d + 2\pi\sqrt{-1}(\xi_1 dy_1 + \xi_2 dy_2).$$

Let $\mathcal{L}$ be the line bundle determined by $\check{\nabla}$. Then the degree of its restriction to E_i is given by

$$\begin{aligned} \deg \mathcal{L}|_{E_i} &= \int_{E_i} \frac{\sqrt{-1}}{2\pi} F_{\check{\nabla}} \\ &= -\int_{E_i} (d\xi_1 \wedge dy_1) \\ &= -(\xi_1(s_i) - \xi_1(s_{i-1})). \end{aligned}$$

We have the second equality because y_2 is constant (and $x_2 = -\lambda = 0$) on E_i. Hence the isomorphism class of the line bundle $\mathcal{L}$ is completely determined by the increment of the angle coordinate ξ_1 on the Lagrangian section L as one moves from $(s_{i-1}, 0)$ to $(s_i, 0)$.

Now we prove Theorem 1.1:

Proof of Theorem 1.1. Recall that the Hamiltonian isotopy class of the Lagrangian submanifold L_γ remains the same when we deform γ in a fixed homotopy class (relative to the ends $\gamma|_{(-\infty,-T]}$ and $\gamma|_{[T,\infty]}$). So, up to a re-parametrization of γ, we can deform the restriction of γ to (s_{i-1}, s_i) to a path arbitrary close to the concatenation of $\gamma_0|_{(s_{i-1},s_i)}$ (the negative axis) with a loop winding around the circle $l_s := \{z \in \mathbb{C}^\times || z| = e^{s_i}\}$ for $w_i(\gamma)$ times. Along γ_0, the angle coordinate ξ_1 is constantly zero, and ξ increases by one when we wind around l_s once in the anti-clockwise direction. Hence, the increment $\xi_1(s_i) - \xi_1(s_{i-1})$ is precisely given by the i-th winding number $w_i(\gamma)$. $\square$

One way to visualize the Lagrangian submanifold L_γ is to observe that the curve $\tau_{s_0}^{s_i}(C_{s_0})$ is given by twisting C_{s_0} in the vanishing cycle for $\sum_{k=1}^{i} w_i(\gamma)$ times. Correspondingly, the increment of the angle coordinate ξ_1 as one goes from $(s_{i-1}, 0)$ to $(s_i, 0)$ is given by the i-th winding number $w_i(\gamma)$ of γ.

5 Wrapped Fukaya categories of suspensions

Let $\mathcal{X}$ be a smooth toric Fano stack of dimension n and $\mathcal{K}$ be the total space of its canonical bundle. The mirror of $\mathcal{X}$ is given by the Lefschetz fibration

$$W : E = (\mathbb{C}^\times)^n \to S = \mathbb{C}$$

defined by a general Laurent polynomial whose Newton polytope coincides with the fan polytope of $\mathcal{X}$. One can choose sufficiently general W so that W is tame (i.e. the gradient $\|\nabla W\|$ is bounded from below by a positive number outside of a compact set) and has only non-degenerate critical points (i.e. the Hessian is non-degenerate around each critical point) with distinct critical values.

There are several ways to associate a Fukaya category to a Lefschetz fibration. One is the *Fukaya-Seidel category* $\mathcal{F}(W)$, which is defined by Seidel [Sei08, Definition 18.12] as the $\mathbb{Z}/2\mathbb{Z}$-invariant part of the Fukaya category of a double cover $\widetilde{E}$ of the total space E branched along the fiber $E_* := W^{-1}(*)$ over the base point. Another one is introduced by Abouzaid [Abo06], which consists of exact Lagrangian submanifolds with boundaries on the fiber E_*. A Lagrangian L in this category must project by W to a curve $\gamma \subset S$ in the neighborhood of the boundary ∂L, and should agree with the parallel transport of ∂L along γ in this neighborhood. The space $\hom(L_1, L_2)$ of morphisms between two objects is defined by perturbing L_2 positively. In this paper, we adopt the following variant of Abouzaid's construction, which follows the original idea of Kontsevich [Kon98] closely:

- For a point $z \in S$, let $\ell_z := \{w \in S \mid \mathfrak{Re}(w) \geq \mathfrak{Re}(z), \mathfrak{Im}(w) = \mathfrak{Im}(z)\}$ be the half line on S starting from z and parallel to the real axis. Assume that $\mathfrak{Re}\, z$ is positive and greater than the real part of any critical value of W. Then for any Lagrangian submanifold L_z of $E_z := W^{-1}(z)$, the trajectory of its parallel transport along ℓ_z gives a Lagrangian submanifold $L \subset E$ fibered over ℓ_z and satisfying $\partial L = L_z$.

- An object of $\mathfrak{Fuk}\, W$ is an *admissible Lagrangian submanifold*, which is an exact Lagrangian submanifold of E which coincides with the Lagrangian submanifold obtained as above for some $z \in S$ and $L_z \subset E_z$ outside of a compact set (see Figure 5.1).

- For a pair (L_1, L_2) of admissible Lagrangian submanifolds, the space $\hom_{\mathfrak{Fuk}\, W}(L_1, L_2)$ of morphisms is defined as the Lagrangian intersection Floer complex $CF^*(L_1^\theta, L_2)$ between L_1^θ and L_2. Here L_1^θ is the Lagrangian submanifold obtained by perturbing the part of L_1 fibered over the half line ℓ_{z_1} to another Lagrangian submanifold fibered over the line $\ell_{z_1}^\theta$ starting from z_1 and extending to the direction of $\exp(\sqrt{-1}\theta)$ for some $0 < \theta \ll 1$ as in Figure 5.2.



Figure 5.1: Admissible Lagrangians



Figure 5.2: Perturbing L_1

- For a sequence $(L_1, \dots, L_k)$ of admissible Lagrangians, the A_∞-operation is defined by perturbing L_i to $L_i^{\theta_i}$ for $0 = \theta_k < \theta_{k-1} < \cdots < \theta_1 \ll 1$ and counting virtual numbers of holomorphic disks bounded by these Lagrangian submanifolds.

- A *vanishing path* is an embedded path $\gamma : [0, \infty) \to S$ which starts from a critical value, avoids the origin and other critical values, and asymptotes to the positive real infinity along a line parallel to the real axis. By arranging the vanishing cycles along a vanishing path, one obtains an admissible Lagrangian submanifold of E called the *Lefschetz thimble.*

- A *distinguished basis of vanishing paths* is a collection $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_m)$ of mutually non-intersecting vanishing paths, one for each critical value and ordered according to the imaginary part of the asymptotic line. The corresponding *distinguished basis of Lefschetz thimbles* will be denoted by $\boldsymbol{\Delta} = (\Delta_1, \dots, \Delta_m)$. The full subcategory of $\mathfrak{Fuk}\, W$ consisting of $\boldsymbol{\Delta}$ will be denoted by $\mathfrak{Fuk}\, \boldsymbol{\Delta}$.

It is expected that $\boldsymbol{\Delta}$ is a full exceptional collection in $D^b \mathfrak{Fuk}\, W$ so that $D^b \mathfrak{Fuk}\, \boldsymbol{\Delta} \cong D^b \mathfrak{Fuk}\, W$ is independent of the choice of $\boldsymbol{\Delta}$, and that $D^b \mathfrak{Fuk}\, W$ is equivalent to the derived Fukaya-Seidel category $D^b \mathcal{F}(W)$. Homological mirror symmetry for toric Fano stacks can be formulated as follows:

Conjecture 5.1 ([Kon98]). *There exists an equivalence*

$$\Psi : D^b \mathfrak{Fuk}\, \boldsymbol{\Delta} \to D^b \operatorname{coh} \mathcal{X} \tag{5.1}$$

of triangulated categories.

Let $\sigma : E \to E$ be a Hamiltonian diffeomorphism which covers the Dehn twist $\overline{\sigma} : S \to S$ along a circle on S with a large radius. The resulting push-forward functor $\sigma_* : \mathfrak{Fuk}\, W \to \mathfrak{Fuk}\, W$ is an autoequivalence of the Fukaya category, which wraps a Lagrangian as shown in Figure 5.3.

Given two admissible Lagrangian submanifolds L_1 and L_2, one can arrange that the Floer complex $\hom_{\mathfrak{Fuk}\, W}(L_1, L_2)$ is a subcomplex of $\hom_{\mathfrak{Fuk}\, W}(\sigma(L_1), L_2)$. This gives a canonical natural transformation

$$\mathfrak{t} : \sigma_* \to \operatorname{id}. \tag{5.2}$$

On the mirror side, there is a distinguished autoequivalence $S(-) = - \otimes \omega_{\mathcal{X}}[n] : D^b \operatorname{coh} \mathcal{X} \to D^b \operatorname{coh} \mathcal{X}$ called the *Serre functor* [BK89]. There is also a distinguished



Figure 5.3: The functor σ_*

section $s \in H^0(\omega_{\mathcal{X}}^\vee)$ characterized by the property that the zero locus $s^{-1}(0)$ is the union of toric divisors of $\mathcal{X}$. This section induces a natural transformation

$$\mathfrak{s} : S[-n] \to \mathrm{id} \tag{5.3}$$

which acts on objects by multiplication by $s : S[-n](X) = X \otimes \omega_{\mathcal{X}} \to \mathrm{id}(X) = X$.

Conjecture 5.2 (Kontsevich, Seidel [Sei09, Example 5]). *The functor σ_* is mirror to the functor $S[-n]$, and the natural transform* (5.2) *is mirror to the natural transformation* (5.3).

The fiber $E_0 := W^{-1}(0)$ at the origin is a divisor in E, which is disjoint from $\sigma^j(\Delta_i)$ for any $i \in \{1, \dots, m\}$ and $j \in \mathbb{Z}$. Hence any holomorphic disk $\varphi : D^2 \to E$ bounded by these Lagrangians has a well-defined intersection number $\varphi(D^2) \cdot E_0$ with E_0. Assume that the collection $(\Delta_1, \dots, \Delta_m)$ is *cyclic*, in the sense that $\mathrm{Ext}^i(\sigma^j(\Delta_k), F_\ell) = 0$ for any $i \neq 0$, any $j \geq 0$ and any $k, \ell \in \{1, \dots, m\}$. Although cyclicity is a strong condition, it is known that any toric Fano stack in dimension two has a cyclic full exceptional collection of line bundles [IU]. It is not known whether any toric Fano stack has a cyclic full exceptional collection of complexes. For $i_1, i_2 \in \{1, \dots, m\}$, $j_1, j_2 \in \mathbb{Z}$ and $x \in \mathrm{Hom}\left((\sigma^{j_1}(\Delta_{i_1}), \sigma^{j_2}(\Delta_{i_2})\right)$, let $\deg(x)$ be the maximal integer d such that x is in the image of

$$\mathfrak{t}^d : \mathrm{Hom}\left(\sigma^{j_1-d}(\Delta_{i_1}), \sigma^{j_2}(\Delta_{i_2})\right) \to \mathrm{Hom}\left(\sigma^{j_1}(\Delta_{i_1}), \sigma^{j_2}(\Delta_{i_2})\right).$$

The following conjecture controls the behavior of the wrapped Fukaya categories under suspension:

Conjecture 5.3. *If a holomorphic map $\varphi : D^2 \to E$ contributes to the x_0-component of $\mathfrak{m}_2(x_1, x_2)$ where $x_0 \in \sigma^{j_0}(\Delta_{i_0}) \cap \sigma^{j_2}(\Delta_{i_2})$, $x_1 \in \sigma^{j_0}(\Delta_{i_0}) \cap \sigma^{j_1}(\Delta_{i_1})$, $x_2 \in \sigma^{j_1}(\Delta_{i_1}) \cap \sigma^{j_2}(\Delta_{i_2})$, and $j_0 \geq j_1 \geq j_2$, then one has $\varphi(D^2) \cdot E_0 = \deg x_0 - \deg x_1 - \deg x_2$.*

The *suspension* of a Lefschetz fibration $W : E \to \mathbb{C}$ is defined by

$$W^\sigma(\boldsymbol{x}, u) = W(\boldsymbol{x}) - u^2 : E^\sigma = E \times \mathbb{C} \to \mathbb{C}.$$

The double suspension is equivalent to

$$W^{\sigma\sigma} = W(\boldsymbol{x}) - uv : E^{\sigma\sigma} = E \times \mathbb{C}^2 \to \mathbb{C}$$

after a change of coordinates. Let

$$Y := \{(\boldsymbol{x}, u, v) \in E \times \mathbb{C}^2 \mid W(\boldsymbol{x}) = uv\}$$

be the fiber of the double suspension at the origin. If $D^b\mathcal{F}(W)$ is equivalent to $D^b \operatorname{coh} \mathcal{X}$, then the double suspension is mirror to the total space of the canonical bundle:

Theorem 5.4 (Seidel [Sei10, Theorem 1]). *There is a full embedding*

$$D^b \operatorname{coh}_0 \mathcal{K} \hookrightarrow D^b \mathfrak{Fuk}\, Y$$

of triangulated categories, where $D^b \operatorname{coh}_0 \mathcal{K}$ is the full subcategory of $D^b \operatorname{coh} \mathcal{K}$ consisting of complexes whose cohomologies are supported on the zero-section.

There is a conic fibration

$$\begin{array}{cccc} \varpi : & Y & \to & E \\ & \rotatebox{90}{$\in$} & & \rotatebox{90}{$\in$} \\ & (\boldsymbol{x}, u, v) & \mapsto & \boldsymbol{x} \end{array}$$

whose discriminant locus is $E_0 := W^{-1}(0)$. Choose a distinguished basis $(\gamma_i)_{i=1}^m$ of vanishing paths as in Figure 5.4, and take the Lagrangian submanifolds $L_i \subset Y$ as in Section 3, which are $\mathbb{R}$-fibrations over the Lefschetz thimbles $\Delta_i \subset E$. Figure 5.5 shows the picture of the Lagrangian on the fiber.



Figure 5.4: Vanishing paths



Figure 5.5: The Lagrangian on the fiber

Choose the wrapping Hamiltonian as the sum $H = H_b + H_f$ of the base Hamiltonian H_b and the fiber Hamiltonian H_f as in [Pas, Section 7]. The base Hamiltonian wraps E in such a way that its image by W is wrapped as shown in Figures 5.6. For the fiber Hamiltonian, there are two choices; $H_{f,1}$ and $H_{f,2}$. The former wraps the Lagrangian around the top of the cylinder while keeping the bottom fixed as shown in Figure 5.7. The latter wraps the Lagrangian on both ends of the cylinder as shown in Figure 5.8

The wrapped Floer cohomology of a pair (L_i, L_j) is given by

$$\operatorname{Hom}_{\mathcal{W}}(L_1, L_2) = \lim_{t \to \infty} \operatorname{Hom}_{\mathfrak{Fuk}\, Y}(\phi_t L_1, L_2)$$

where $\phi_t : M \to M$ is the flow generated by the Hamiltonian H. The cochain complex underlying the wrapped Floer cohomology is obtained by telescope construction, and the



Figure 5.6: Wrapping the base



Figure 5.7: Wrapping the fiber by $H_{f,1}$



Figure 5.8: Wrapping the fiber by $H_{f,2}$

A_∞-operations are defined by using *cascades*, which combine the A_∞-operations and the continuation maps in Floer theory [AS10, Abo10, Aur10]. We write the A_∞-categories consisting of $\{L_i\}_{i=1}^m$ and wrapped by $H_1 = F_b + H_{f,1}$ and $H_2 = F_b + F_{f,2}$ as $\mathcal{W}_1$ and $\mathcal{W}_2$ respectively.

Theorem 5.5. *Assume that the collection* $\mathbf{\Delta}$ *is cyclic. Then Conjectures 5.1, 5.2 and 5.3 imply equivalences*

$$D^b\mathcal{W}_1 \cong D^b \operatorname{coh} \mathcal{K} \tag{5.4}$$

and

$$D^b\mathcal{W}_2 \cong D^b \operatorname{coh} \check{\mathcal{Y}} \tag{5.5}$$

of triangulated categories.

Proof. We first show the equivalence (5.4). Note that the monodromy of the conic fibration $\varpi : Y \to E$ around the discriminant $E_0 = W^{-1}(0)$ is given by the Dehn twist along the vanishing cycle, which is inverse to the wrapping on the fiber. It follows that for a positive integer t, the intersection points $\phi_t(L_i) \cap L_j$ are bijective with $\coprod_{k=0}^t \left(\sigma^k(\Delta_i) \cap \Delta_j\right)$, so that one has

$$\operatorname{Hom}_{\mathfrak{Fuk}\, Y}(\phi_t(L_i), L_j) = \bigoplus_{k=0}^{t} \operatorname{Hom}_{\mathfrak{Fuk}\, W}(\sigma^k(\Delta_i), \Delta_j), \tag{5.6}$$

cf. Figures 5.7, 5.9 and 5.10. It follows that there is an isomorphism



Figure 5.9: Wrapping once



Figure 5.10: Wrapping twice

$$\begin{aligned}
\mathrm{Hom}_{\mathcal{W}_1}(L_i, L_j) &= \lim_{t\to\infty} \mathrm{Hom}_{\mathfrak{Fuk}\, Y}(\phi_t(L_i), L_j) \\
&\cong \bigoplus_{k=0}^{\infty} \mathrm{Hom}_{\mathfrak{Fuk}\, W}(\sigma^k(\Delta_i), \Delta_j) \\
&\cong \bigoplus_{k=0}^{\infty} \mathrm{Hom}_{\mathcal{X}}(E_i \otimes \omega_{\mathcal{X}}^k, E_j) \\
&\cong \bigoplus_{k=0}^{\infty} \mathrm{Hom}_{\mathcal{X}}(E_i, E_j \otimes \omega_{\mathcal{X}}^{-k}) \\
&\cong \mathrm{Hom}_{\mathcal{X}}\left(E_i, \bigoplus_{k=0}^{\infty} E_j \otimes \omega_{\mathcal{X}}^{-k}\right) \\
&\cong \mathrm{Hom}_{\mathcal{X}}(E_i, E_j \otimes \pi_* \mathcal{O}_{\mathcal{K}}) \\
&\cong \mathrm{Hom}_{\mathcal{X}}(E_i, \pi_* \pi^* E_j) \\
&\cong \mathrm{Hom}_{\mathcal{K}}(\pi^* E_i, \pi^* E_j)
\end{aligned}$$

of $\mathbb{C}$-vector spaces, where $\pi : \mathcal{K} \to \mathcal{X}$ is the natural projection. We take the isomorphism

$$\bigoplus_{k=0}^{\infty} \mathrm{Hom}_{\mathfrak{Fuk}\, W}(\sigma^k(\Delta_i), \Delta_j) \cong \bigoplus_{k=0}^{\infty} \mathrm{Hom}_{\mathcal{X}}(E_i \otimes \omega_{\mathcal{X}}^k, E_j)$$

in such a way that an element

$$x \in \mathrm{Hom}_{\mathfrak{Fuk}\, W}(\sigma^k(\Delta_i), \Delta_j)$$

is sent to

$$(s-1)^d y' \in \bigoplus_{\ell=0}^{d} \mathrm{Hom}_{\mathcal{X}}(E_i \otimes \omega_X^{k-\ell}, E_j),$$

where $d = \deg x$, $x = \mathfrak{t}^d(x')$ and the element $y' \in \mathrm{Hom}_{\mathcal{X}}(E_i \otimes \omega_X^{k-d}, E_j)$ is mirror to $x' \in \mathrm{Hom}_{\mathfrak{Fuk}\, W}(\sigma^{k-d}\Delta_i, \Delta_j)$ under homological mirror symmetry for $\mathcal{X}$.

Any holomorphic map $\varphi : D^2 \to Y$ bounded by $\phi_t(L_i)$'s is a holomorphic section of the conic fibration $\varpi : Y \to E$ over the image $\varpi \circ \varphi(D^2) \subset S$ of the holomorphic map $\varpi \circ \varphi : D^2 \to E$ bounded by $\sigma^k(\Delta_i)$'s. This gives the relation between holomorphic disks in E contributing to compositions in $\mathfrak{Fuk}\, W$ and holomorphic disks in Y contributing to $\mathfrak{Fuk}\, Y$. The precise formula for this contribution is calculated by Pascaleff [Pas, Proposition 4.4] as

$$\mu^2_{\mathfrak{Fuk}\, Y}\left(\mathfrak{t}^{d_1}(x_1'), \mathfrak{t}^{d_2}(x_2')\right) = \sum_{s=0}^{k} \binom{k}{s} \mathfrak{t}^{d_1+d_2+s} x_3',$$

where $k = \deg(\mu^2_{\mathfrak{Fuk}\, W}(x_1', x_2'))$ and $\mu^2_{\mathfrak{Fuk}\, W}(x_1', x_2') = \mathfrak{t}^k(x_3')$. Note that the integer k is equal to the intersection number of the disk contributing to $\mu^2_{\mathfrak{Fuk}\, W}(x_1', x_2')$ by Conjecture 5.3. This matches the composition

$$\begin{aligned}(s-1)^{d_1} y_1' \cdot (s-1)^{d_2} y_2' &= (s-1)^{d_1+d_2} y_1' y_2' \\ &= (s-1)^{d_1+d_2} s^k y_3' \\ &= \sum_{t=0}^{k} \binom{k}{t} (s-1)^{d_1+d_2+t} y_3'\end{aligned}$$

in $D^b \operatorname{coh} \mathcal{K}$, where $k = \deg(y_1' y_2')$ and $y_1' y_2' = s^k y_3'$.

Since $\mathbf{\Delta}$ is cyclic, all the morphisms have degree zero, so that A_∞-operations μ^k for $k \neq 2$ is trivial. Since the collection $(E_1, \dots, E_m)$ is full, its pull-back $(\pi^* E_1, \dots, \pi^* E_m)$ generates $D^b \operatorname{coh} \mathcal{K}$, and the equivalence

$$D^b \mathcal{W}_1 \cong D^b \operatorname{coh} \mathcal{K}$$

is proved.

If one uses H_2 as the wrapping Hamiltonian, then one obtains

$$\operatorname{Hom}_{\mathfrak{Fuk}\, Y}(\phi_t(L_i), L_j) = \bigoplus_{k=0}^{2t} \operatorname{Hom}_{\mathfrak{Fuk}\, W}(\sigma^k(\Delta_i), \Delta_j), \tag{5.7}$$

instead of (5.6), and the continuation map

$$\operatorname{Hom}_{\mathfrak{Fuk}\, Y}(\phi_t(L_i), L_j) \to \operatorname{Hom}_{\mathfrak{Fuk}\, Y}(\phi_{t+1}(L_i), L_j)$$

is mirrored by the inclusion

$$\bigoplus_{k=0}^{2t} \operatorname{Hom}_{\mathcal{X}}(E_i \otimes \omega_{\mathcal{X}}^k, E_j) \hookrightarrow \bigoplus_{k=0}^{2t+1} \operatorname{Hom}_{\mathcal{X}}(E_i \otimes \omega_{\mathcal{X}}^k, E_j)$$

followed by multiplication by $s-1$;

$$s-1 : \bigoplus_{k=0}^{2t+1} \operatorname{Hom}_{\mathcal{X}}(E_i \otimes \omega_{\mathcal{X}}^k, E_j) \to \bigoplus_{k=0}^{2t+2} \operatorname{Hom}_{\mathcal{X}}(E_i \otimes \omega_{\mathcal{X}}^k, E_j).$$

The resulting direct limit gives the space of morphisms on the complement $\check{\mathcal{Y}} \subset \mathcal{K}$ of the zero locus of $s-1$, and the equivalence (5.5) is proved. $\square$

6 Minimal resolution of A_n-singularity

Let p and q be positive integers satisfying $n+1 = p+q$ and

$$W = z^p + \frac{1}{z^q} : \mathbb{C}^\times \to \mathbb{C}$$

be the Landau-Ginzburg potential for the mirror of the orbifold curve $\mathbb{X}_{p,q}$ with two orbifold points of weights p and q. The fiber $W^{-1}(R)$ of W at a sufficiently large positive real number R consists of $p+q$ points, p of which are close to p-th roots of R and q of which are close to q-th roots of $1/R$. It follows that an admissible Lagrangian submanifold on the torus $\mathbb{C}^\times = \operatorname{Spec} \mathbb{C}[z, z^{-1}]$ asymptotes to $\zeta_p^i \cdot \mathbb{R}^+$ for some i near $z \to \infty$ and $\zeta_q^j \cdot \mathbb{R}^+$ for some j near $z \to 0$, where we put $\zeta_p = \exp(2\pi\sqrt{-1}/p)$ and $\zeta_q = \exp(2\pi\sqrt{-1}/q)$.

Now set $(p,q) = (1,n)$ so that $W = z + 1/z^n$ and $\mathbb{X}_{1,n} \cong \mathbb{P}(1,n)$ is a weighted projective line. It suffices to work out this case since the total space $\mathcal{K} = \mathcal{K}_{\mathbb{X}_{p,q}}$ of the canonical bundle is derived equivalent to the minimal resolution of the A_n-singularity $\mathbb{C}^2/(\mathbb{Z}/n\mathbb{Z})$ for any pair (p,q) satisfying $p+q = n+1$. This is related to the function $1/z + z^n$ appearing in Introduction by an automorphism $z \mapsto 1/z$ of the torus, and our choice in this section is made only for aesthetic reason (Figure 6.2 looks nicer for this choice). The critical points of W are given by $z = \sqrt[n+1]{n}\zeta_{n+1}^i$, $i = 0, \dots, n$, with critical values $\sqrt[n+1]{n}(1+1/n)\zeta_{n+1}^i$. The fiber $W^{-1}(R)$ for a large positive real number R consists of one point which is approximately R and n points which are approximately n-th roots of $1/R$.

Take the distinguished basis $(\gamma_i)_{i=0}^n$ of vanishing paths as in Figure 6.1. The corresponding Lefschetz thimbles look as in Figure 6.2. Let $\gamma_{-(n+1)k+i}$ be the vanishing path obtained by applying $\overline{\sigma}^k$ to γ_i for $i = \{0, \dots, n\}$ and $k \in \mathbb{Z}$. The corresponding Lefschetz thimble will be denoted by $\Delta_{-(n+1)k+i} = \sigma^k(\Delta_i)$.



Figure 6.1: Lefschetz thimbles on the W-plane



Figure 6.2: Lefschetz thimbles on the z-plane

The direct sum $\bigoplus_{i,j=0}^{m}\bigoplus_{k=0}^{\infty}\operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^k(\Delta_i),\Delta_j)$ has a ring structure given by

$$\begin{array}{ccccc}
\operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^{k_2}(\Delta_{i_2}),\Delta_{i_3}) & \otimes & \operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^{k_1}(\Delta_{i_1}),\Delta_{i_2}) & \to & \operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^{k_1+k_2}(\Delta_{i_1}),\Delta_{i_3}) \\
\rotatebox{90}{$\in$} & & \rotatebox{90}{$\in$} & & \rotatebox{90}{$\in$} \\
x_2 & & x_1 & \mapsto & \mu^2_{\mathfrak{Fuk}\,W}(x_2,\sigma_*^{k_2}(x_1))
\end{array}$$

on components. The natural transformation (5.2) induces an endomorphism $\mathfrak{T}$ of this ring defined by

$$\begin{array}{ccc}
\operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^{k}(\Delta_{i}),\Delta_{j}) & \to & \operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^{k+1}(\Delta_{i}),\Delta_{j}) \\
\rotatebox{90}{$\in$} & & \rotatebox{90}{$\in$} \\
x & \mapsto & \mu^2_{\mathfrak{Fuk}\,W}(x,\mathfrak{t}(\sigma^k(\Delta_i)))
\end{array}$$

on components.

Let $\mathbb{C}[x,y]$ be the homogeneous coordinate ring of $\mathcal{X}=\mathbb{P}(1,n)$, where $\deg x=1$ and $\deg y=n$. The canonical bundle is given by $\omega_{\mathcal{X}}=\mathcal{O}(-n-1)$, and the exceptional collection

$$(E_0,E_1,\dots,E_n)=(\mathcal{O},\mathcal{O}(1),\dots,\mathcal{O}(n))$$

of line bundles is full and cyclic. The ring structure on $\bigoplus_{i,j=0}^{m}\bigoplus_{k=0}^{\infty}\operatorname{Hom}_{\mathcal{X}}(E_i\otimes\omega_{\mathcal{X}}^k,E_j)$ is defined similarly as above, and the ring endomorphism $\mathfrak{S}$ induced by the natural transformation (5.3) is the multiplication by $xy\in H^0(\omega_{\mathcal{X}}^{\vee})$.

The following proposition gives the parts of Conjectures 5.1 and 5.2 which are needed for the proof of Theorem 5.5:

Proposition 6.1. *There is a ring isomorphism*

$$\bigoplus_{i,j=0}^{m}\bigoplus_{k=0}^{\infty}\operatorname{Hom}_{\mathfrak{Fuk}\,W}(\sigma^k(\Delta_i),\Delta_j)\xrightarrow{\sim}\bigoplus_{i,j=0}^{m}\bigoplus_{k=0}^{\infty}\operatorname{Hom}_{\mathcal{X}}(E_i\otimes\omega_{\mathcal{X}}^k,E_j)\tag{6.1}$$

which sends the ring endomorphism $\mathfrak{T}$ *to* $\mathfrak{S}$.

Proof. The Lefschetz thimble $\Delta_{-k(n+1)+i}$ starts from the critical point $\sqrt[n+1]{n}\,\zeta_{n+1}^i$ and extends in two directions: One wraps around the origin k/n times and approaches the origin in the direction of ζ_n^{i-k}. The other wraps around infinity k times and asymptotes to the real line. Figure 6.3 shows the picture of two Lefschetz thimbles. Here the top and the bottom edges of the rectangle are identified to form a cylinder, which is obtained as the real blow-up of $\mathbb{P}^1$ at the origin and infinity. The vertical dotted line shows the locus where the absolute value is $\sqrt[n+1]{n}$. The Lefschetz thimble Δ_0 is just the real line, and the Lefschetz thimble $\Delta_{-k(n+1)+i}$ is obtained from the Lefschetz thimble Δ_i by wrapping k times.

Write $k(n+1)-i=k'n+i'$ for $k'\in\mathbb{Z}$ and $i'\in\{0,\dots,n-1\}$. Then the thimbles $\Delta_{-k(n+1)+i}$ and Δ_0 intersect at $k'+1$ points, and we label these intersection points by the basis of $\operatorname{Hom}(\mathcal{O}_{\mathbb{P}(1,n)}(-k(n+1)+i),\mathcal{O}_{\mathbb{P}(1,n)})$ as shown in Figure 6.3. Intersections between other Lefschetz thimbles can be labeled similarly. One can easily arrange that $\operatorname{Hom}_{\mathfrak{Fuk}\,W}(\Delta_i,\Delta_j)$ for $i\le j$ have degree zero (and those for $i>j$ have degree one), so that there are no higher A_∞-operations on $\bigoplus_{i,j=0}^{m}\bigoplus_{k=0}^{\infty}\hom_{\mathfrak{Fuk}\,W}(\sigma^k(\Delta_i),\Delta_j)$. It is also easy to see that the composition $\mu^2_{\mathfrak{Fuk}\,W}$ on the left hand side of (6.1) (coming from triangles on $\mathbb{C}^\times$) matches the composition on the right hand side (which is just the multiplication of polynomials), and Proposition 6.1 is proved. $\square$



Figure 6.3: Intersections of Lefschetz thimbles

The proposition below is an analogue of [Pas, Proposition 4.3], which provides the part of Conjecture 5.3 needed for the proof of Theorem 5.5:

Proposition 6.2. *If a triangle D on $\mathbb{C}^\times$ contributing to the composition on the left hand side of* (6.1) *corresponds to the multiplication*

$$x^{i_1}y^{j_1} \cdot x^{i_2}y^{j_2} = x^{i_3}y^{j_3}$$

on the right hand side of (6.1)*, then one has*

$$D \cdot W^{-1}(0) = d_3 - d_1 - d_2, \tag{6.2}$$

where $D \cdot W^{-1}(0)$ is the intersection number of the triangle with the divisor $W^{-1}(0) \subset \mathbb{C}^\times$, and d_ℓ for $\ell = 1, 2, 3$ is the number of times $x^{i_\ell}y^{j_\ell}$ can be divided by $x^n y$.

Proof. Since $\sqrt[n+1]{n}$ is close to 1, one can perturb Figure 6.3 slightly to set the dotted vertical line to be the unit circle on $\mathbb{C}^\times$ without changing the intersections of triangles with $W^{-1}(0)$. Then $W^{-1}(0)$ are equidistributed on the dotted vertical line with vertical coordinates $(2j+1)\pi/(2n+2)$ for $j = 0, \dots, n$. Let L_1, L_2 and L_3 be the Lagrangians bounding D, so that $x^{i_1}y^{j_1}$, $x^{i_2}y^{j_2}$ and $x^{i_1}y^{j_1}$ label a point in $L_1 \cap L_2$, $L_2 \cap L_3$ and $L_1 \cap L_3$ respectively. The triangle looks either as in Figure 6.4 or Figure 6.5.



Figure 6.4: A triangle



Figure 6.5: Another triangle

Write $x^{i_\ell}y^{j_\ell} = (xy)^{d_\ell}x^{i_\ell - d_\ell}y^{j_\ell - d_\ell}$ for $\ell = 1, 2, 3$. If $i_\ell - d_\ell = 0$ for $\ell = 1, 2$ or $j_\ell - d_\ell = 0$ for $\ell = 1, 2$, then the whole triangle is either on the left or on the right of the dotted vertical line in Figure 6.3, and both sides of (6.2) is zero. If $a := i_1 - d_1 > 0$ and $b := j_2 - d_2 > 0$, then the triangle D looks as in Figure 6.4, and the vertical dotted line cuts D into two. The number of points in $W^{-1}(0)$ on the part of the vertical dotted line



Figure 6.6: The dual cycles

bounded by L_1 and L_2 is given by a, and that by L_2 and L_3 is given by b (here we are working on the universal cover of the cylinder $\mathbb{C}^\times$). The vertical dotted line is on the right or the left of the vertex $x^{i_3}y^{j_3}$ depending on whether $a > b$ or $a < b$ (and exactly on the line if $a = b$). It follows that both sides of (6.2) is given by $\max\{a, b\}$ in this case. The case $j_1 - d_1 > 0$ and $i_2 - d_2 > 0$ is similar, and Proposition 6.2 is proved. □

This concludes the proof of Theorem 1.2.

Now we discuss the compatibility of Theorem 1.1 and Theorem 1.2. Let $(\epsilon_1, \dots, \epsilon_n)$ be the collection of line segments on E connecting points in $W^{-1}(0)$ as in Figure 6.6, so that the intersection numbers with Lefschetz thimbles are given by

$$\Delta_i \cdot \epsilon_j = \delta_{ij}, \quad i = 0, \dots, n, \ j = 1, \dots, n.$$

One can choose a symplectomorphism $Y \xrightarrow{\sim} Y'$ in such a way that these intersection numbers correspond precisely to the winding numbers in Definition 3.2. The SYZ transformations of the resulting Lagrangians in Y are then given by line bundles $(\mathcal{L}_i)_{i=0}^n$ on $\check{Y}$ satisfying

$$\deg \mathcal{L}_i|_{E_j} = \delta_{ij}$$

for $i = 0, \dots, n$ and $j = 1, \dots, n$.

On the other hand, the proof of the equivalence $D^b\mathcal{W}_2 \cong D^b \operatorname{coh} \mathcal{Y}$ in this section (note that the A_∞ category $\mathcal{W}_2$ here is denoted by $\mathcal{W}'$ in (1.7)) gives a collection $(\mathcal{L}_i')_{i=0}^n$ of line bundles on $\check{\mathcal{Y}}$ defined by

$$\mathcal{L}_i' = \pi^* \mathcal{O}_{\mathbb{P}(1,n)}(i)\big|_{\check{\mathcal{Y}}}.$$

One can easily see that by sending $\mathcal{L}_i'$ to $\mathcal{L}_i$ for $i = 0, \dots, n$, one obtains an equivalence $D^b \operatorname{coh} \check{\mathcal{Y}} \xrightarrow{\sim} D^b \operatorname{coh} \check{Y}$. This gives the compatibility of Theorem 1.1 and Theorem 1.2 discussed in Introduction.

References

[AAK] Mohammed Abouzaid, Denis Auroux, and Ludmil Katzarkov, *Lagrangian fibrations on blowups of toric varieties and mirror symmetry for hypersurfaces*, arXiv:1205.0053.

[Abo06] Mohammed Abouzaid, *Homogeneous coordinate rings and mirror symmetry for toric varieties*, Geom. Topol. **10** (2006), 1097–1157 (electronic). MR MR2240909 (2007h:14052)

[Abo09] ______, *Morse homology, tropical geometry, and homological mirror symmetry for toric varieties*, Selecta Math. (N.S.) **15** (2009), no. 2, 189–270. MR MR2529936

[Abo10] ______, *A geometric criterion for generating the Fukaya category*, Publ. Math. Inst. Hautes Études Sci. (2010), no. 112, 191–240. MR 2737980

[AP01] D. Arinkin and A. Polishchuk, *Fukaya category and Fourier transform*, Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), AMS/IP Stud. Adv. Math., vol. 23, Amer. Math. Soc., Providence, RI, 2001, pp. 261–274. MR 1876073 (2002k:32040)

[AS10] Mohammed Abouzaid and Paul Seidel, *An open string analogue of Viterbo functoriality*, Geom. Topol. **14** (2010), no. 2, 627–718. MR 2602848 (2011g:53190)

[Aur07] Denis Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. GGT **1** (2007), 51–91. MR 2386535 (2009f:53141)

[Aur09] ______, *Special Lagrangian fibrations, wall-crossing, and mirror symmetry*, Surveys in differential geometry. Vol. XIII. Geometry, analysis, and algebraic geometry: forty years of the Journal of Differential Geometry, Surv. Differ. Geom., vol. 13, Int. Press, Somerville, MA, 2009, pp. 1–47. MR MR2537081

[Aur10] ______, *Fukaya categories of symmetric products and bordered Heegaard-Floer homology*, J. Gökova Geom. Topol. GGT **4** (2010), 1–54. MR 2755992

[BK89] A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1183–1205, 1337. MR MR1039961 (91b:14013)

[Cha] Kwokwai Chan, *Homological mirror symmetry for A_n-resolutions as a T-duality*, to appear in J. London Math. Soc., arXiv:1112.0844.

[Cha10] ______, *The Ooguri-Vafa metric, holomorphic discs and wall-crossing*, Math. Res. Lett. **17** (2010), no. 3, 401–414. MR 2653678 (2011g:53187)

[CLL12] K. Chan, S.-C. Lau, and N.C. Leung, *SYZ mirror symmetry for toric Calabi-Yau manifolds*, J. Diff. Geom. **90** (2012), no. 2, 177–250.

[Dui80] J. J. Duistermaat, *On global action-angle coordinates*, Comm. Pure Appl. Math. **33** (1980), no. 6, 687–706. MR 596430 (82d:58029)

[Fan08] Bohan Fang, *Homological mirror symmetry is T-duality for $\mathbb{P}^n$*, Commun. Number Theory Phys. **2** (2008), no. 4, 719–742. MR MR2492197

[FLTZ11] Bohan Fang, Chiu-Chu Melissa Liu, David Treumann, and Eric Zaslow, *The coherent-constructible correspondence and homological mirror symmetry for toric varieties*, Geometry and analysis. No. 2, Adv. Lect. Math. (ALM), vol. 18, Int. Press, Somerville, MA, 2011, pp. 3–37. MR 2882439

[FLTZ12] ______, *T-duality and homological mirror symmetry for toric varieties*, Adv. Math. **229** (2012), no. 3, 1875–1911. MR 2871160 (2012m:14075)

[Fuk05] Kenji Fukaya, *Multivalued Morse theory, asymptotic analysis and mirror symmetry*, Graphs and patterns in mathematics and theoretical physics, Proc. Sympos. Pure Math., vol. 73, Amer. Math. Soc., Providence, RI, 2005, pp. 205–278. MR 2131017 (2006a:53100)

[GS11] Mark Gross and Bernd Siebert, *From real affine geometry to complex geometry*, Ann. of Math. (2) **174** (2011), no. 3, 1301–1428. MR 2846484

[Hit01] Nigel Hitchin, *Lectures on special Lagrangian submanifolds*, Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), AMS/IP Stud. Adv. Math., vol. 23, Amer. Math. Soc., Providence, RI, 2001, pp. 151–182. MR 1876068 (2003f:53086)

[IU] Akira Ishii and Kazushi Ueda, *Dimer models and exceptional collections*, arXiv:0911.4529.

[Kon95] Maxim Kontsevich, *Homological algebra of mirror symmetry*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 120–139. MR MR1403918 (97f:32040)

[Kon98] ______, *Lectures at ENS Paris, spring 1998*, set of notes taken by J. Bellaiche, J.-F. Dat, I. Martin, G. Rachinet and H. Randriambololona, 1998.

[KS06a] Maxim Kontsevich and Yan Soibelman, *Affine structures and non-Archimedean analytic spaces*, The unity of mathematics, Progr. Math., vol. 244, Birkhäuser Boston, Boston, MA, 2006, pp. 321–385. MR MR2181810

[KS06b] ______, *Notes on A_∞ algebras, A_∞ categories and non-commutative geometry. I*, arXiv:math/0606241, 2006.

[KV00] M. Kapranov and E. Vasserot, *Kleinian singularities, derived categories and Hall algebras*, Math. Ann. **316** (2000), no. 3, 565–576. MR MR1752785 (2001h:14012)

[LYZ00] Naichung Conan Leung, Shing-Tung Yau, and Eric Zaslow, *From special Lagrangian to Hermitian-Yang-Mills via Fourier-Mukai transform*, Adv. Theor. Math. Phys. **4** (2000), no. 6, 1319–1341. MR 1894858 (2003b:53053)

[Pas] James Pascaleff, *Floer cohomology in the mirror of the projective plane and a binodal cubic curve*, arXiv:1109.3255.

[Sei08] Paul Seidel, *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR MR2441780

[Sei09] ______, *Symplectic homology as Hochschild homology*, Algebraic geometry—Seattle 2005. Part 1, Proc. Sympos. Pure Math., vol. 80, Amer. Math. Soc., Providence, RI, 2009, pp. 415–434. MR 2483942 (2010c:53129)

[Sei10] ______, *Suspending Lefschetz fibrations, with an application to local mirror symmetry*, Comm. Math. Phys. **297** (2010), no. 2, 515–528. MR 2651908

[SYZ96] Andrew Strominger, Shing-Tung Yau, and Eric Zaslow, *Mirror symmetry is T-duality*, Nuclear Phys. B **479** (1996), no. 1-2, 243–259. MR MR1429831 (97j:32022)

Kwokwai Chan
Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong
e-mail address : kwchan@math.cuhk.edu.hk

Kazushi Ueda
Department of Mathematics, Graduate School of Science, Osaka University, Machikaneyama 1-1, Toyonaka, Osaka, 560-0043, Japan.
e-mail address : kazushi@math.sci.osaka-u.ac.jp